arXiv:1301.6432v1 [math.CA] 28 Jan 2013

A NEW PROOF OF THE GEOMETRIC-ARITHMETIC MEAN INEQUALITY BY CAUCHY'S INTEGRAL FORMULA

FENG QI, XIAO-JING ZHANG, AND WEN-HUI LI

Abstract. Let $a=(a_1,a_2,\dots,a_n)$ for $n\in\mathbb{N}$ be a given sequence of positive numbers. In the paper, the authors establish, by using Cauchy's integral formula in the theory of complex functions, an integral representation of the principal branch of the geometric mean

$$G_n(a+z)=\left[\prod_{k=1}^n(a_k+z)\right]^{1/n}$$

for $z\in\mathbb{C}\setminus(-\infty,-\min\{a_k,1\le k\le n\}]$, and then provide a new proof of the well known GA mean inequality.

1. Introduction

Let $a=(a_1,a_2,\dots,a_n)$ for $n\in\mathbb{N}$, the set of all positive integers, be a given sequence of positive numbers. Then the arithmetic and geometric means $A_n(a)$ and $G_n(a)$ of the numbers $a_1,a_2,\dots,a_n$ are defined respectively as

$$A_n(a)=\frac{1}{n}\sum_{k=1}^n a_k \tag{1.1}$$

and

$$G_n(a)=\left(\prod_{k=1}^n a_k\right)^{1/n}. \tag{1.2}$$

It is common knowledge that

$$G_n(a)\le A_n(a), \tag{1.3}$$

with equality if and only if $a_1=a_2=\dots=a_n$.

There has been a large number, presumably over one hundred, of proofs of the GA mean inequality (1.3) in the mathematical literature. The most complete information, so far, can be found in the monographs [1, 2, 3, 4, 5, 6] and a lot of references therein.

In this paper, we establish, by using Cauchy's integral formula in the theory of complex functions, an integral representation of the principal branch of the geometric mean

$$G_n(a+z)=\left[\prod_{k=1}^n(a_k+z)\right]^{1/n} \tag{1.4}$$

2010 *Mathematics Subject Classification.* Primary 26E60, 30E20; Secondary 26A48, 44A20.

Key words and phrases. Integral representation; Cauchy's integral formula; Arithmetic mean; Geometric mean; GA mean inequality; New proof.

This paper was typeset using $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-LaTeX.

for $z\in\mathbb{C}\setminus(-\infty,-\min\{a_k,1\le k\le n\}]$, and provide a new proof of the GA mean inequality (1.3).

2. Lemmas

In order to prove our main results, we need the following lemmas.

Lemma 2.1. *For $z\in\mathbb{C}\setminus(-\infty,-\min\{a_k,1\le k\le n\}]$, the principal branch of the complex function*

$$f_n(z)=G_n(a+z)-z, \tag{2.1}$$

where $a+z=(a_1+z,a_2+z,\dots,a_n+z)$, meets

$$\lim_{z\to\infty}f_n(z)=A_n(a). \tag{2.2}$$

Proof. By L'Hôspital's rule in the theory of complex functions, we have

$$\begin{aligned}\lim_{z\to\infty}f_n(z)&=\lim_{z\to\infty}\left\{z\left[G_n\left(1+\frac{a}{z}\right)-1\right]\right\}\\&=\lim_{z\to0}\frac{G_n(1+az)-1}{z}=\lim_{z\to0}\frac{\mathrm{d}}{\mathrm{d}z}\left[\prod_{k=1}^{n}(1+a_kz)\right]^{1/n}=A_n(a),\end{aligned}$$

where $1+\frac{a}{z}=\left(1+\frac{a_1}{z},1+\frac{a_2}{z},\dots,1+\frac{a_n}{z}\right)$ and $1+az=(1+a_1z,1+a_2z,\dots,1+a_nz)$. Lemma 2.1 is thus proved. □

Lemma 2.2. *For $z\in\mathbb{C}\setminus(-\infty,0]$ and $a=(a_1,a_2,\dots,a_n)$ satisfying $a_\ell\le a_{\ell+1}$ for $1\le\ell\le n-1$, let*

$$h_n(z)=G_n(a-a_1+z)-z, \tag{2.3}$$

where $a-a_1+z=(z,a_2-a_1+z,\dots,a_n-a_1+z)$. Then the imaginary part of the principal branch of $h_n(z)$ meets

$$\lim_{\varepsilon\to0^+}\Im h_n(-t+i\varepsilon)=\begin{cases}\left[\prod_{k=1}^{n}|a_k-a_1-t|\right]^{1/n}\sin\dfrac{\ell\pi}{n}, & t\in(a_\ell-a_1,a_{\ell+1}-a_1]\\0, & t>a_n-a_1\end{cases} \tag{2.4}$$

for $1\le\ell\le n-1$.

Proof. For $t=a_{\ell+1}-a_1$ for $1\le\ell\le n-1$, we have

$$\begin{aligned}h_n(-t+i\varepsilon)&=\exp\left[\frac{1}{n}\sum_{k\ne\ell+1}^{n}\ln(a_k-a_1-t+i\varepsilon)+\frac{1}{n}\ln(i\varepsilon)\right]+t-i\varepsilon\\&=\exp\left[\frac{1}{n}\sum_{k\ne\ell+1}^{n}\ln(a_k-a_1-t+i\varepsilon)\right]\exp\left[\frac{1}{n}\left(\ln|\varepsilon|+\frac{\pi}{2}i\right)\right]+t-i\varepsilon\\&\to\exp\left[\frac{1}{n}\sum_{k\ne\ell+1}^{n}\ln(a_k-a_1-t)\right]\lim_{\varepsilon\to0^+}\exp\left[\frac{1}{n}\left(\ln|\varepsilon|+\frac{\pi}{2}i\right)\right]+t\\&=t\end{aligned}$$

as $\varepsilon\to0^+$. Hence, when $t=a_{\ell+1}-a_1$ for $1\le\ell\le n-1$, we have

$$\lim_{\varepsilon\to0^+}\Im h_n(-t+i\varepsilon)=0.$$

For $t \in (0, \infty) \setminus \{a_{\ell+1} - a_1, 1 \le \ell \le n-1\}$ and $\varepsilon > 0$, we have

$$\begin{aligned}
h_n(-t+i\varepsilon) &= G_n(a - a_1 - t + i\varepsilon) + t - i\varepsilon \\
&= \exp\left[\frac{1}{n}\sum_{k=1}^{n} \ln(a_k - a_1 - t + i\varepsilon)\right] + t - i\varepsilon \\
&= \exp\left\{\frac{1}{n}\sum_{k=1}^{n} [\ln|a_k - a_1 - t + i\varepsilon| + i\arg(a_k - a_1 - t + i\varepsilon)]\right\} + t - i\varepsilon \\
&\to \begin{cases} \exp\left(\dfrac{1}{n}\displaystyle\sum_{k=1}^{n} \ln|a_k - a_1 - t| + \dfrac{\ell\pi}{n} i\right) + t, & t \in (a_\ell - a_1, a_{\ell+1} - a_1) \\ \exp\left(\dfrac{1}{n}\displaystyle\sum_{k=1}^{n} \ln|a_k - a_1 - t| + \pi i\right) + t, & t > a_n - a_1 \end{cases} \\
= &\begin{cases} \left(\displaystyle\prod_{k=1}^{n} |a_k - a_1 - t|\right)^{1/n} \left(\cos\dfrac{\ell\pi}{n} + i\sin\dfrac{\ell\pi}{n}\right) + t, & t \in (a_\ell - a_1, a_{\ell+1} - a_1) \\ \left(\displaystyle\prod_{k=1}^{n} |a_k - a_1 - t|\right)^{1/n} (\cos\pi + i\sin\pi) + t, & t > a_n - a_1 \end{cases}
\end{aligned}$$

as $\varepsilon \to 0^+$. As a result, we have

$$\lim_{\varepsilon \to 0^+} \Im h_n(-t+i\varepsilon) = \begin{cases} \left(\displaystyle\prod_{k=1}^{n} |a_k - a_1 - t|\right)^{1/n} \sin\dfrac{\ell\pi}{n}, & t \in (a_\ell - a_1, a_{\ell+1} - a_1); \\ 0, & t > a_n - a_1. \end{cases}$$

The proof of Lemma 2.2 is completed. $\square$

3. An integral representation of the geometric mean

Now we are in a position to establish an integral representation of the geometric mean $G_n(a+z)$.

Theorem 3.1. *Let $0 < a_k \le a_{k+1}$ for $1 \le k \le n-1$ and $a + z = (a_1 + z, a_2 + z, \dots, a_n + z)$ for $z \in \mathbb{C} \setminus (-\infty, -a_1]$. Then the principal branch of the geometric mean $G_n(a+z)$ has the integral representation*

$$G_n(a+z) = A_n(a) + z - \frac{1}{\pi}\sum_{\ell=1}^{n-1} \sin\frac{\ell\pi}{n} \int_{a_\ell}^{a_{\ell+1}} \left|\prod_{k=1}^{n} (a_k - t)\right|^{1/n} \frac{\mathrm{d}t}{t+z}. \tag{3.1}$$

Proof. By standard arguments, it is not difficult to see that

$$\lim_{z \to 0^+} [z h_n(z)] = 0 \quad \text{and} \quad h_n(\overline{z}) = \overline{h_n(z)}. \tag{3.2}$$

For any but fixed point $z \in \mathbb{C} \setminus (-\infty, 0]$, choose $0 < \varepsilon < 1$ and $r > 0$ such that $0 < \varepsilon < |z| < r$, and consider the positively oriented contour $C(\varepsilon, r)$ in $\mathbb{C} \setminus (-\infty, 0]$ consisting of the half circle $z = \varepsilon e^{i\theta}$ for $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and the half lines $z = x \pm i\varepsilon$ for $x \le 0$ until they cut the circle $|z| = r$, which close the contour at the points $-r(\varepsilon) \pm i\varepsilon$, where $0 < r(\varepsilon) \to r$ as $\varepsilon \to 0$. By the famous Cauchy's integral formula in the theory of complex functions, we have

$$\begin{aligned}
h_n(z)&=\frac{1}{2\pi i}\oint_{C(\varepsilon,r)}\frac{h_n(w)}{w-z}\,\mathrm{d}w\\
&=\frac{1}{2\pi i}\biggl[\int_{\pi/2}^{-\pi/2}\frac{i\varepsilon e^{i\theta}h\bigl(\varepsilon e^{i\theta}\bigr)}{\varepsilon e^{i\theta}-z}\,\mathrm{d}\theta+\int_{\arg[-r(\varepsilon)-i\varepsilon]}^{\arg[-r(\varepsilon)+i\varepsilon]}\frac{ire^{i\theta}h\bigl(re^{i\theta}\bigr)}{re^{i\theta}-z}\,\mathrm{d}\theta\\
&\quad+\int_{-r(\varepsilon)}^{0}\frac{h_n(x+i\varepsilon)}{x+i\varepsilon-z}\,\mathrm{d}x+\int_{0}^{-r(\varepsilon)}\frac{h_n(x-i\varepsilon)}{x-i\varepsilon-z}\,\mathrm{d}x\biggr].
\end{aligned}\tag{3.3}$$

By the limit in (3.2), it follows that

$$\lim_{\varepsilon\to0^+}\int_{\pi/2}^{-\pi/2}\frac{i\varepsilon e^{i\theta}h_n\bigl(\varepsilon e^{i\theta}\bigr)}{\varepsilon e^{i\theta}-z}\,\mathrm{d}\theta=0.\tag{3.4}$$

By virtue of the limit (2.2) in Lemma 2.1, we deduce that

$$\begin{aligned}
\lim_{\substack{\varepsilon\to0^+\\ r\to\infty}}\int_{\arg[-r(\varepsilon)-i\varepsilon]}^{\arg[-r(\varepsilon)+i\varepsilon]}\frac{ire^{i\theta}h_n\bigl(re^{i\theta}\bigr)}{re^{i\theta}-z}\,\mathrm{d}\theta&=\lim_{r\to\infty}\int_{-\pi}^{\pi}\frac{ire^{i\theta}h_n\bigl(re^{i\theta}\bigr)}{re^{i\theta}-z}\,\mathrm{d}\theta\\
&=2A_n(a-a_1)\pi i,
\end{aligned}\tag{3.5}$$

where $a-a_1=(0,a_2-a_1,\dots,a_n-a_1)$. Utilizing the second formula in (3.2) and the limit (2.4) in Lemma 2.2 results in

$$\begin{aligned}
&\int_{-r(\varepsilon)}^{0}\frac{h_n(x+i\varepsilon)}{x+i\varepsilon-z}\,\mathrm{d}x+\int_{0}^{-r(\varepsilon)}\frac{h_n(x-i\varepsilon)}{x-i\varepsilon-z}\,\mathrm{d}x\\
&=\int_{-r(\varepsilon)}^{0}\biggl[\frac{h_n(x+i\varepsilon)}{x+i\varepsilon-z}-\frac{h_n(x-i\varepsilon)}{x-i\varepsilon-z}\biggr]\,\mathrm{d}x\\
&=\int_{-r(\varepsilon)}^{0}\frac{(x-i\varepsilon-z)h_n(x+i\varepsilon)-(x+i\varepsilon-z)h_n(x-i\varepsilon)}{(x+i\varepsilon-z)(x-i\varepsilon-z)}\,\mathrm{d}x\\
&=\int_{-r(\varepsilon)}^{0}\frac{(x-z)[h_n(x+i\varepsilon)-h_n(x-i\varepsilon)]-i\varepsilon[h_n(x-i\varepsilon)+h_n(x+i\varepsilon)]}{(x+i\varepsilon-z)(x-i\varepsilon-z)}\,\mathrm{d}x\\
&=2i\int_{-r(\varepsilon)}^{0}\frac{(x-z)\Im h_n(x+i\varepsilon)-\varepsilon\Re h_n(x+i\varepsilon)}{(x+i\varepsilon-z)(x-i\varepsilon-z)}\,\mathrm{d}x\\
&\to2i\int_{-r}^{0}\frac{\lim_{\varepsilon\to0^+}\Im h_n(x+i\varepsilon)}{x-z}\,\mathrm{d}x\\
&=-2i\int_{0}^{r}\frac{\lim_{\varepsilon\to0^+}\Im h_n(-t+i\varepsilon)}{t+z}\,\mathrm{d}t\\
&\to-2i\int_{0}^{\infty}\frac{\lim_{\varepsilon\to0^+}\Im h_n(-t+i\varepsilon)}{t+z}\,\mathrm{d}t\\
&=-2i\sum_{\ell=1}^{n-1}\sin\frac{\ell\pi}{n}\int_{a_\ell-a_1}^{a_{\ell+1}-a_1}\Biggl[\prod_{k=1}^{n}|a_k-a_1-t|\Biggr]^{1/n}\frac{\mathrm{d}t}{t+z}
\end{aligned}\tag{3.6}$$

as $\varepsilon \to 0^+$ and $r \to \infty$. Substituting equations (3.4), (3.5), and (3.6) into (3.3) and simplifying generate

$$h_n(z) = A_n(a - a_1) - \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin \frac{\ell\pi}{n} \int_{a_\ell - a_1}^{a_{\ell+1} - a_1} \left[\prod_{k=1}^{n} |a_k - a_1 - t| \right]^{1/n} \frac{\mathrm{d}t}{t + z}. \tag{3.7}$$

From $f_n(z) = h_n(z + a_1) + a_1$ and (3.7), it is immediate to deduce that

$$\begin{aligned} f_n(z) &= A_n(a - a_1) + a_1 - \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin \frac{\ell\pi}{n} \int_{a_\ell - a_1}^{a_{\ell+1} - a_1} \left[\prod_{k=1}^{n} |a_k - a_1 - t| \right]^{1/n} \frac{\mathrm{d}t}{t + z + a_1} \\ &= A_n(a) - \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin \frac{\ell\pi}{n} \int_{a_\ell}^{a_{\ell+1}} \left[\prod_{k=1}^{n} |a_k - t| \right]^{1/n} \frac{\mathrm{d}t}{t + z}, \end{aligned}$$

from which the integral representation (3.1) follows. Theorem 3.1 is proved. □

4. A NEW PROOF OF THE GA MEAN INEQUALITY

With the aid of the integral representation (3.1) in Theorem 3.1, we can easily deduce the GA mean inequality (1.3) as follows.

Taking $z = 0$ in the integral representation (3.1) yields

$$G_n(a) = A_n(a) - \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin \frac{\ell\pi}{n} \int_{a_\ell}^{a_{\ell+1}} \left[\prod_{k=1}^{n} |a_k - t| \right]^{1/n} \frac{\mathrm{d}t}{t} \le A_n(a).$$

From this, it is also obvious that the equality in (1.3) is valid if and only if $a_1 = a_2 = \cdots = a_n$. The proof of the GA mean inequality (1.3) is complete.

REFERENCES

[1] E. F. Beckenbach and R. Bellman, *Inequalities*, Springer, Berlin, 1983.

[2] P. S. Bullen, *Handbook of Means and Their Inequalities*, Mathematics and its Applications (Dordrecht) **560**, Kluwer Academic Publishers, Dordrecht, 2003.

[3] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, 2nd ed., Cambridge University Press, Cambridge, 1952.

[4] J.-C. Kuang, *Chángyòng Bùděngshì* (*Applied Inequalities*), 3rd ed., Shāndōng Kēxué Jìshù Chūbǎn Shè (Shandong Science and Technology Press), Ji'nan City, Shandong Province, China, 2004. (Chinese)

[5] D. S. Mitrinović, *Analytic Inequalities*, Springer, New York-Heidelberg-Berlin, 1970.

[6] D. S. Mitrinović and P. M. Vasić, *Sredine*, Matematička Biblioteka **40**, Beograd, 1969.

(Qi) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA
E-mail address: `qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com`
URL: `http://qifeng618.wordpress.com`

(Zhang) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA
E-mail address: `xiao.jing.zhang@qq.com`

(Li) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA
E-mail address: `wen.hui.li@foxmail.com`